

Could 2S 0114+650 be a magnetar?

X.-D. Li^{1,2} and E. P. J. van den Heuvel²

ABSTRACT

We investigate the spin evolution of the binary X-ray pulsar 2S 0114+650, which possesses the slowest known spin period of ~ 2.7 hours. We argue that, to interpret such long spin period, the magnetic field strength of this pulsar must be initially $\gtrsim 10^{14}$ G, that is, it was born as a magnetar. Since the pulsar currently has a normal magnetic field $\sim 10^{12}$ G, our results present support for magnetic field decay predicted by the magnetar model.

Subject headings: binaries: close - pulsars: individual: 2S 0114+650 - stars: neutron - X-ray: stars

1. Introduction

Neutron stars are thought to be born as rapidly rotating (~ 10 ms) radio pulsars created during a type II/Ib supernova explosion involving a massive star. The dipolar magnetic fields of radio pulsars, as inferred from their observed spin-down rates, range from 10^8 G to 3×10^{13} G (Taylor, Manchester, & Lyne 1993). However, it has been proposed that there may exist “magnetars” - neutron stars with magnetic field strengths in excess of $\sim 10^{14}$ G (Thompson & Duncan 1992). These objects have been invoked to model soft gamma-ray repeaters (SGRs) and the 6 – 12 s anomalous X-ray pulsars (AXPs) (Thompson & Duncan 1996; see also Kouveliotou et al. 1993, 1994; van Paradijs, Taam, & van den Heuvel 1995; Corbet et al. 1995; Vasisht, Frail, & Kulkarni 1995; Vasisht & Gotthelf 1997), though unambiguous evidence for the existence of magnetars comes from recent observations of the AXP-like object 1E 1841–045 (Vasisht & Gotthelf 1997), SGRs 1806–20 (Kouveliotou et al. 1998a) and 1900+14 (Kouveliotou et al. 1998b).

According to Thompson & Duncan (1992, 1996), magnetars are neutron stars born with millisecond periods that generate magnetic fields above 10^{14} G by dynamo action due to convective turbulence, magnetic field decay powers the X-ray and particle emission of magnetars. It is conceivable that when the magnetic fields in magnetars have decayed to, say, 10^{12} G, and accretion occurs, they are not distinguished from X-ray pulsars born with normal ($10^{12} - 10^{13}$ G) magnetic fields, except that they may have relatively longer spin periods. One may then expect to find magnetar descendants among long-period binary X-ray pulsars, because they are much more luminous in X-rays than isolated objects accreting from interstellar medium. Here we present arguments indicating that 2S 0114+650, the X-ray pulsar with the slowest known spin, may have a magnetar evolutionary history.

2. The slowest X-ray pulsar 2S 0114+650

The X-ray source 2S 0114+650 was discovered in 1977 by the SAS 3 galactic survey (Dower & Kelly 1977). Its optical counterpart, LS I+65 010, was recently identified as a supergiant of spectral type B1 Ia

¹ Department of Astronomy, Nanjing University, Nanjing 210093, China

² Astronomical Institute, University of Amsterdam, Kruislaan 403, 1098 SJ Amsterdam, The Netherlands

(Reig et al. 1996). Thus 2S 0114+650 belongs to the class of high-mass X-ray binaries (HMXBs), systems in which a compact star - generally a neutron star - accompanies a high-mass donor star (cf. Bhattacharya & van den Heuvel 1991 for a review). With a distance of 7.2 kpc derived from this spectral classification, the X-ray luminosity is a few $10^{35} - 10^{36} \text{ erg s}^{-1}$. An orbital period of 11.59 days was reported from optical radial velocity measurements by Crampton, Hutchings, & Cowley (1985). There has been some weak evidence of a pulsation period of 850 – 895 s (Yamauchi et al. 1990; Koenigsberger et al. 1983). In contrast, Finley, Belloni, & Cassinelli (1992) have discovered periodic X-ray outbursts with a 2.78 hour period. The same period was confirmed by ROSAT observations (Finley, Taylor, & Belloni 1994). Recent Rossi X-ray Timing Explorer (RXTE) observations show presence of modulation on 11.59 day orbital period as well as 2.7 hour pulse period (Corbet, Finley, & Peele 1998). If this pulse period indeed represents the rotation period of the neutron star, 2S 0114+650 would be by far the slowest known X-ray pulsar.

3. The spin evolution in 2S 0114+650

How has 2S 0114+650 been spun down to the long period ($P \simeq 10^4 \text{ s}$) if it was formed with much shorter period ($\sim 0.01 - 1 \text{ s}$, say)? The neutron star’s spin evolution is divided in three phases (see Davies & Pringle 1981; Bhattacharya & van den Heuvel 1991). In the first radio pulsar phase, the star is an active radio pulsar and spins down by magnetic dipole radiation. This phase ends when the ram pressure of the ambient material overcomes the pulsar’s wind pressure at the gravitational radius ($R_G = 2GM/V^2$, where G is the gravitation constant, M is the neutron star mass, and V is the relative velocity between the neutron star and the ambient material). In the second propeller phase material enters the corotating magnetosphere and is stopped at R_A , the Alfvén radius, where the energy density in the accretion flow balances the local magnetic pressure. Further penetration cannot occur owing to the centrifugal barrier; that is, $R_A > R_c$, where $R_c = [GM(P/2\pi)^2]^{1/3}$ is the corotation radius. The star expels the material once it spins up the material to the local escape velocity at $\sim R_A$ (Illarionov & Sunyaev 1975). Propeller action continues until $R_A \simeq R_c$, when the centrifugal barrier is removed and the spin period reaches its equilibrium value (Bhattacharya & van den Heuvel 1991)

$$P_{\text{eq}} \simeq (20 \text{ s}) B_{12}^{6/7} \dot{M}_{15}^{-3/7} R_6^{18/7} M_{1.4}^{-5/7}, \quad (1)$$

where $B = 10^{12} B_{12} \text{ G}$ is the neutron star’s dipolar magnetic field strength, $\dot{M} = 10^{15} \dot{M}_{15} \text{ g s}^{-1}$ the mass accretion rate, $R = 10^6 R_6 \text{ cm}$ the radius, and $M_{1.4} = M/1.4 M_\odot$ (Throughout this Letter, we take $M_{1.4} = R_6 = 1$). In the following accretion phase the star becomes an X-ray pulsar, and its spin evolution is determined by the net torque exerted on the star by the accretion flow.

The X-ray emission observed in 2S 0114+650 is most likely to be powered by accretion onto the neutron star via a stellar wind from the companion. With a mass-loss rate of a few $10^{-6} M_\odot \text{ yr}^{-1}$ and a wind velocity of $\sim 10^3 \text{ km s}^{-1}$, the derived X-ray luminosity from a simple wind-fed model is in accordance with the mean detected one (Reig et al. 1996). A stable accretion disk is unlikely to form around the neutron star, which would require an extremely low ($\sim 200 \text{ km s}^{-1}$) wind velocity (Wang 1981). Early two-dimensional numerical studies of Bondi-Hoyle accretion flow (e.g., Matsuda, Inoue, & Swada 1987; Fryxell & Taam 1988, 1989) demonstrated that temporary accretion disks with alternating sense of rotation possibly form in a wind accreting system. More recent high resolution three-dimensional numerical investigations (e.g., Ruffert 1992, 1997) found the so-called wind “flip-flop” instability with the timescale of the order of hours. This is consistent with the torque fluctuations in wind-accreting X-ray pulsars (Nagase 1989), suggesting that the long-term averaged angular momentum transferred to the neutron star by the accreted wind material is

very small. We therefore conclude that the spin period of 2S 0114+650 has not been considerably changed during the present accretion phase. This means that the long spin period of 2S 0114+650 was actually attained in an earlier evolutionary phase before the companion star became a super-giant - it was spun down by the propeller mechanism (Illarionov & Sunyaev 1975) when the companion star was on main-sequence and had a weaker wind. The magnitude of the equilibrium period, as seen in equation (1), is determined by the magnetic field strength and mass accretion rate of the neutron star during this phase.

The magnetic field strength in 2S 0114+650 has not been measured, since no cyclotron features have been seen in its X-ray spectrum. But there is indirect evidence indicating that its magnetic field strength is similar to those in other X-ray pulsars: Its spectrum shows the typical shape of the usual X-ray pulsars having a power-law with an exponential high-energy cutoff at ~ 7 keV (Yamauchi et al. 1990) or ≥ 15 keV (Koenigsberger et al. 1983). The iron emission line at about 6.4 keV, which is common among X-ray pulsars, was also discovered in 2S 0114+650 (Yamauchi et al. 1990; Apparao, Bisht, & Singh 1991). Observations of cyclotron lines in X-ray pulsars imply surface fields of about $(0.5 - 5) \times 10^{12}$ G, and it has been suggested that the cutoff seen in the power-law spectra of X-ray pulsars is related to the magnetic field strength of the neutron star (Makishima & Mihara 1992)³. This would imply a field of $\sim 10^{12}$ G for 2S 0114+650, consistent with those for typical X-ray pulsars which show cutoff energies of 10 – 20 keV.

The mass of LS I+65 010 was estimated to be $16(\pm 5)M_{\odot}$ from evolutionary models (Reig et al. 1996). For a typical mass-loss rate of $\sim 10^{-8} - 10^{-7} M_{\odot} \text{ yr}^{-1}$ from a $16 M_{\odot}$ main-sequence star and a wind velocity of $\sim 10^3 \text{ km s}^{-1}$, the accretion rate of the neutron star is $\dot{M} \sim 10^{13} - 10^{14} \text{ g s}^{-1}$. Inserting the values of B and $\dot{M}$ into equation (1), we find $P_{\text{eq}} \sim 50 - 140$ s, nearly two orders of magnitude shorter than the observed period. (Even if B is enhanced to 10^{13} G, P_{eq} never exceeds ~ 1000 s.) This implies that the extremely long period of 2S 0114+650 cannot be reached via the propeller mechanism *if the pulsar has possessed a constant magnetic field of $\sim 10^{12} - 10^{13}$ G*. One may argue that the accretion rate during the propeller phase could be much lower than that adopted here, because of a lower rate of mass loss from the companion star or a higher wind velocity. For example, if $\dot{M}$ ranges from $\sim 10^9 \text{ g s}^{-1}$ (for $B = 10^{12}$ G) to $\sim 10^{11} \text{ g s}^{-1}$ (for $B = 10^{13}$ G), the value of P_{eq} can be indeed raised to $\sim 10^4$ s. However, this would lead to a spin-down time during the radio pulsar phase (Davies & Pringle 1981)

$$\begin{aligned} \tau_s &\simeq (2.5 \times 10^{10} \text{ yr}) B_{12}^{-1} \dot{M}_9^{-1/2} V_8^{-1} \\ &\simeq (2.5 \times 10^8 \text{ yr}) B_{13}^{-1} \dot{M}_{11}^{-1/2} V_8^{-1} \end{aligned} \quad (2)$$

(where $V = 10^8 V_8 \text{ cm s}^{-1}$), which is much longer than the lifetime the companion star spends on main-sequence (generally $\lesssim 10^7 \text{ yr}$).

There exists another possibility that the neutron star was born rotating slowly ($P \sim 1$ s), so that it went directly to the propeller phase. Propeller spin-down to P_{eq} takes (Wang & Robertson 1985)

$$\begin{aligned} \tau_s &\simeq (1.5 \times 10^{10} \text{ yr}) B_{12}^{-1/2} \dot{M}_9^{-3/4} P_1^{-3/4} \\ &\simeq (1.5 \times 10^8 \text{ yr}) B_{13}^{-1/2} \dot{M}_{11}^{-3/4} P_1^{-3/4} \end{aligned} \quad (3)$$

(where $P_1 = P/1$ s), which is still much longer than 10^7 yr .

We are eventually led to the conclusion that 2S 0114+650 must initially have had a magnetic field much stronger than its present value, that is, it was born as a magnetar. Distinguished from neutron

³This relation is not quite accurate (see Reynolds, Parmar, & White 1993), but it may provide an order of magnitude estimate of the magnetic fields.

stars with normal magnetic field strengths, magnetars are radio-quiet, and the decaying magnetic fields power the X-ray and particle emission. For a magnetar in a binary system, during the early evolutionary phase, the pressure exerted by particle emission exceeds the stellar wind ram pressure at $\sim R_G$, preventing accretion onto the neutron star (Thompson & Duncan 1996). After $t \simeq 10^4 - 10^5$ years, the star's spin period increases to

$$P \simeq (10 \text{ s}) t_4^{1/2} B_{15} R_6^2 M_{1.4}^{-1/2} \quad (4)$$

by magnetic dipole radiation (where $t_4 = t/10^4$ yr), the particle luminosity decays, and the wind material from the companion star begins to interact with the magnetosphere. Again adopt a mass accretion rate of $\sim 10^{13} - 10^{14} \text{ g s}^{-1}$, the propeller effect, for a magnetic field of $\sim (1 - 4) \times 10^{14} \text{ G}$, can comfortably spin down the neutron star from $\sim 10 \text{ s}$ to $\sim 10^4 \text{ s}$ on a timescale $\lesssim 10^5$ years (cf. equations [1] and [3]), i.e., before the magnetic field decays significantly (see below). After this long equilibrium period is reached, the period of the neutron star would remain close to it, due to inefficient angular momentum transfer during the subsequent wind-accretion process.

Field decay in 2S 0114+650 should be slow enough to guarantee the final, long spin period, and fast enough to allow a considerable reduction in the field within the lifetime of the system. There exist several mechanisms for field decay in non-accreting neutron stars: Ohmic decay, ambipolar diffusion and Hall drift (e.g., Goldreich & Reisenegger 1992). Ohmic decay dominates in weakly magnetized ($B \lesssim 10^{11} \text{ G}$) neutron stars, fields in intermediate strength ($B \sim 10^{12} - 10^{13} \text{ G}$) decay via Hall drift, and intense fields ($B \gtrsim 10^{14} \text{ G}$) are mainly affected by ambipolar diffusion. For typical values of the characteristic length scale (10^5 cm) of the flux loops through the outer core and the core temperature (10^8 K), the field-decay timescale for ambipolar diffusion through the solenoidal mode in a magnetar is around $3 \times 10^5 \text{ yr}$ if $B \sim 10^{14} \text{ G}$ (Thompson & Duncan 1996). Compared to the age (a few 10^6 yr) of the optical companion of 2S 0114+650, this implies that the magnetar has had enough time for its field to decay to its current value (a few 10^{12} G). Similar conclusion can also be found from the detailed calculations by Heyl & Kulkarni (1998).

4. Discussion

SGRs, which are transient gamma-ray sources that undergo repeated outbursts, have been suggested to be the prototypes of magnetars. There are four known SGRs, with two (SGRs 1806–20 and 0525–66) associated with young supernova remnants (Kulkarni & Frail 1993; Vasisht et al. 1994). Recently, pulsations at a period of 7.47 s and 5.16 s with a spin-down rate of $2.6 \times 10^{-3} \text{ s yr}^{-1}$ and $3.5 \times 10^{-3} \text{ s yr}^{-1}$ were discovered in the persistent X-ray flux of SGRs 1806–20 (Kouveliotou et al. 1998a) and 1900+14 (Kouveliotou et al. 1998b), respectively; the magnetic field strengths of $(2 - 8) \times 10^{14} \text{ G}$ can be derived if the spin-down is due to magnetic dipole radiation. AXPs are another type of high-energy sources that have recently joined this group of highly magnetized neutron stars. They are a group of about eight pulsating X-ray sources with periods around 6–12 s (cf. Stella, Israel, & Mereghetti 1998 for a review), characterized by steady spin-down, relatively low and constant X-ray luminosities of $\sim 10^{35} - 10^{36} \text{ erg s}^{-1}$, and very soft X-ray spectra. So far no optical counterpart has been detected. Nearly half of them are located at the centers of supernova remnants, suggesting that they are relatively young ($\lesssim 10^5$ years). Dipole magnetic fields of $10^{14} - 10^{15} \text{ G}$ can also be derived from the measured spin-down rates, if they are spinning down due to dipole radiation torques. The observed X-ray luminosities, spin periods and spin-down rates in SGRs and AXPs follow naturally from the magnetar model (Thompson & Duncan 1996).

A crucial test of the magnetar model should include the observational evidence of the hypothesized magnetic field decay in magnetars. This evidence is most likely to be found in slowly-rotating, binary (such

as 2S 0114+650) and isolated (such as RX J0720.4–3125; Haberl et al. 1997; Heyl & Hernquist 1998; Kulkarni & van Kerkwijk 1998) X-ray pulsars. Figure 1 compares the magnetar candidates (SGRs 1806–20 and 1900+14 in filled rectangles, and AXPs 1E 2259+586, 4U 0142+61, 1E 1048.1–5937 and 1E 1841–05 in filled stars) with their possible X-ray pulsar descendants (2S 0114+650 in open triangle and other pulsars in open stars) in the magnetic field versus spin period diagram. A schematic view of the magnetar field and spin evolution of magnetars is clearly seen. The magnetic fields of SGRs and AXPs are derived from their observed spin-down rates (Kouveliotou et al. 1998a, 1998b; Stella, Israel, & Mereghetti 1998 and references therein) under the assumption that the spin-down is caused by magnetic dipole radiation. For 2S 0114+650 we take a canonical magnetic field of 3×10^{12} G with the possible range of $10^{12} - 10^{13}$ G. The dashed lines represent the relation between the magnetic field strength and the equilibrium spin period with various mass accretion rates (10^{18} , 10^{15} and 10^{12} g s^{-1}), from which an initial magnetic field of a pulsar can be constrained by the present spin period, given a reasonable estimate of the mass accretion rate during the propeller phase. The detailed evolutionary track of a magnetar depends on the particle luminosity, the properties of the stellar wind from the companion, the magnetic field and its decay timescale.




Fig. 1.— The magnetic field versus spin period diagram for the candidates of magnetars (SGRs in filled rectangles and AXPs in filled stars) and the slow X-ray pulsars (in open stars except 2S 0114+650 in open triangle) with known magnetic fields. The dashed lines denote the relation between the magnetic field and the equilibrium spin period with various mass accretion rates.

In Fig. 1 we have also plotted the slow (spin periods $\gtrsim 100$ s) X-ray pulsars (A 0535+26, Vela X-1, GX 1+4, 4U 1907+09, 4U 1538–52 and GX 301–2) with known magnetic field strengths as possible descendants of magnetars. Their magnetic field strengths are generally estimated from the observed cyclotron line features in X-ray spectra (Mihara & Makishima 1998 and references therein). For GX 1+4, it is determined from the observational evidence of the propeller effect (Cui 1997). These sources may have had a similar evolutionary history as 2S 0114+650, but for them the evidence is not as unequivocal as for 2S 0114+650. Their slow spins could be accounted for in terms of the propeller effects with normal magnetic field strengths (e.g., Illarionov & Sunyaev 1975; Waters & van Kerkwijk 1989, see also the dashed lines). However, a magnetar model presents an alternative explanation that can not be ruled out. Actually, high magnetic field strengths ($\gtrsim 10^{13}$ G) have been measured in A 0535+26 and GX 1+4. Statistically, if the birth rate of magnetars is of the order of one per millennium (Kouveliotou et al. 1998a), and the X-ray lifetime of HMXBs lasts up to $\sim 10^5$ yr, there may exist in the Galaxy ~ 10 binary X-ray pulsars

originating from magnetars.

The authors thank Jan van Paradijs and the referee for helpful comments. This work was in part supported by National Natural Science Foundation of China and by the Netherlands Organization for Scientific Research (NWO) through Spinoza Grant 08-0 to E. P. J. van den Heuvel.

REFERENCES

- Apparao, K. M. V., Bisht, P., & Singh, K. P. 1991, *ApJ*, 371, 772
- Bhattacharya, D. & van den Heuvel, E. P. J. 1991, *Phys. Rep.*, 203, 1
- Corbet, R. H. D., Finley, P., & Peele, A. G. 1998, *ApJ*, submitted (astro-ph/9809216)
- Corbet, R. H. D., Smale, A. P., Ozaki, M., Koyama, K., & Iwasawa, K. 1995, *ApJ*, 443, 786
- Crampton, D., Hutchings, J., & Cowley, A. 1985, *ApJ*, 299, 839
- Cui, W. 1997, *ApJ*, 482, L163
- Davies, R. E. & Pringle, J. E. 1981, *MNRAS*, 196, 209
- Dower, R. & Kelly, R. 1977, *IAU Circ. No.* 3144
- Finley, J., Belloni, T., & Cassinelli, I. P. 1992, *A&A*, 262, L25
- Finley, J., Taylor, M., & Belloni, T. 1994, *ApJ*, 429, 356
- Fryxell, B. E. & Taam, R. E. 1988, *ApJ*, 335, 862
- . 1989, *ApJ*, 339, 297
- Goldreich, P. & Reisenegger, A. 1992, *ApJ*, 395, 250
- Haberl, F., Motch, C., Buckley, D. A. H., Zickgraf, F. J., & Pietsch, W. 1997, *A&A*, 326, 662
- Heyl, J. & Hernquist, L. 1998, *MNRAS*, 297, L69
- Heyl, J. S. & Kulkarni, S. R. 1998, *ApJ*, 506, L61
- Illarionov, A. F. & Sunyaev, R. A. 1975, *A&A*, 39, 185
- Koenigsberger, G., Swank, J. H., Szymkowiak, A. E., & White, N. E. 1983, *ApJ*, 268, 782
- Kouveliotou, C., Dieters, S., Strohmayer, T., van Paradijs, J., Fishman, G. J. et al. 1998a, *Nature*, 393, 235
- Kouveliotou, C., Fishman, G. J., Meegan, C. A., Paciesas, W. S., van Paradijs, J. et al. 1994, *Nature*, 368, 125
- Kouveliotou, C., Fishman, G. J., Meegan, C. A., Paciesas, W. S., Wilson, R. B. et al. 1993, *Nature*, 362, 728
- Kouveliotou, C., Strohmayer, T., Hurley, K., van Paradijs, J., Finger, M. H. et al. 1998b, *ApJ*, submitted (astro-ph/9809140)

- Kulkarni, S. R. & Frail, D. A. 1993, *Nature*, 365, 33
- Kulkarni, S. R. & van Kerkwijk, M. H. 1998, *ApJ*, 507, L49
- Makishima, K. & Mihara, T. 1992, in *Frontiers in X-ray Astronomy*, ed. Tanaka, Y. & Koyama, K. (Uni. Acad. Press, Tokyo), 23
- Matsuda, T., Inoue, M., & Swada, K. 1987, *MNRAS*, 226, 785
- Mihara, T. & Makishima, K. 1998, in *Neutron Stars and Pulsars*, ed. Shibasaki, N., Kawai, N., Shibata, S., & Kifune, T. (Uni. Acad. Press, Tokyo), 205
- Nagase, F. 1989, *PASJ*, 41, 1
- Reig, P., Chakrabarty, D., Coe, M. J., Fabregat, J., Negueruela, I. et al. 1996, *A&A*, 311, 879
- Reynolds, A. P., Parmar, A. N., & White, N. E. 1993, *ApJ*, 414, 302
- Ruffert, M. 1992, *A&A*, 265, 82
- . 1997, *A&A*, 317, 793
- Stella, L., Israel, G. L., & Mereghetti, S. 1998, in *The Many Faces of Neutron Stars*, ed. Bucceri, R., van Paradijs, J., & Alpar, M. A. (Kluwer Academic Publishers), 397
- Taylor, J. H., Manchester, R. N., & Lyne, A. G. 1993, *ApJS*, 88, 529
- Thompson, C. & Duncan, R. 1992, *ApJ*, 392, L9
- . 1996, *ApJ*, 473, 322
- van Paradijs, J., Taam, R. E., & van den Heuvel, E. P. J. 1995, *A&A*, 299, L41
- Vasisht, G., Frail, D. A., & Kulkarni, S. R. 1995, *ApJ*, 440, L65
- Vasisht, G. & Gotthelf, E. V. 1997, *ApJ*, 486, L129
- Vasisht, G., Kulkarni, S. R., Frail, D. A., & Greiner, J. 1994, *ApJ*, 431, L35
- Wang, Y.-M. 1981, *A&A*, 102, 36
- Wang, Y.-M. & Robertson, J. A. 1985, *A&A*, 151, 361
- Waters, L. F. B. M. & van Kerkwijk, M. H. 1989, *A&A*, 223, 196
- Yamauchi, S., Asaoka, I., Kawada, M., Koyama, K., & Tawara, Y. 1990, *PASJ*, 42, L53